arXiv

Multi-scale Analysis for Rosseland Equation with Small Periodic Oscillating Coefficients

Zhang Qiao-fu

Academy of Mathematics and Systems Science,
Chinese Academy of Sciences

NNSFC(No. 90916027) May, 2012

具有小周期振荡系数的 Rosseland方程多尺度分析

张乔夫

中国科学院数学与系统科学研究院

毕业答辩—2012.05.29

国家自然科学基金(90916027)

Outline

1. Rosseland Eq.—fixed point
    - Model
    - Elliptic
    - Parabolic
2. Math algorithms
3. SOTS—homogenization
4. Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$
    - Correctors estimates
    - Error estimates

Outline

1. Rosseland Eq.—fixed point
    - Model
    - Elliptic
    - Parabolic
2. Math algorithms
3. SOTS—homogenization
4. Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$
    - Correctors estimates
    - Error estimates

Outline

1. Rosseland Eq.—fixed point
    - Model
    - Elliptic
    - Parabolic
2. Math algorithms
3. SOTS—homogenization
4. Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$
    - Correctors estimates
    - Error estimates

Outline

1. Rosseland Eq.—fixed point
   - Model
   - Elliptic
   - Parabolic
2. Math algorithms
3. SOTS—homogenization
4. Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$
   - Correctors estimates
   - Error estimates

Background (1)


zsy-002 10.0kV 14.5mm x2.00k SE(U)
20.0um
zsy-003 10.0kV 14.3mm x2.00k SE(U)
20.0um

SEM of fibrous (H.I.T. 2009)

Nonsmooth data

Background (1)


zsy-002 10.0kV 14.5mm x2.00k SE(U)
20.0um
zsy-003 10.0kV 14.3mm x2.00k SE(U)
20.0um

SEM of fibrous (H.I.T. 2009)
Nonsmooth data

Background (2)



Coat of a rocket (L.I.R.R., 2005).

Background (3)



Temperature (K) (L.I.R.R., 2005).

Background (4)



No blow-up, maximum principles, safe !

Background (4)



No blow-up, maximum principles, safe !

Background (4)



No blow-up, maximum principles, safe !

Part 1: existence / elliptic

Optically thick, Rosseland Eq.

$$-\mathrm{div}[A\nabla u] = f. \tag{1}$$

u: temperature. K, B are S P D.

$$A(u,x) = K(x) + 4\,u^3\,B(x). \tag{2}$$

Conductive, radiative.

Part 1: existence / elliptic

Optically thick, Rosseland Eq.

$$-\mathrm{div}[A\nabla u] = f. \tag{1}$$

u: temperature. K, B are S P D.

$$A(u,x) = K(x) + 4\,u^3\,B(x). \tag{2}$$

Conductive, radiative.

Part 1: existence / elliptic

Optically thick, Rosseland Eq.

$$-\mathrm{div}[A\nabla u] = f. \tag{1}$$

u: temperature. K, B are S P D.

$$A(u,x) = K(x) + 4\,u^3\,B(x). \tag{2}$$

Conductive, radiative.

Part 1: existence / elliptic

Optically thick, Rosseland Eq.

$$-\mathrm{div}[A\nabla u] = f. \tag{1}$$

u: temperature. K, B are S P D.

$$A(u,x) = K(x) + 4\,u^3\, B(x). \tag{2}$$

Conductive, radiative.

Part 1: existence / elliptic

Optically thick, Rosseland Eq.

$$-\mathrm{div}[A\nabla u] = f. \tag{1}$$

u: temperature. K, B are S P D.

$$A(u,x) = K(x) + 4\,u^3\,B(x). \tag{2}$$

Conductive, radiative.

Exists ?

Laitinen, 2002



Remark 3.4. The existence theory for diffusion approximations is not entirely trivial. If conduction were not present the question could be resolved using the well-known Kirchoff transform [8,15]. In the general case more sophisticated analysis in needed. For example the methods of [10] could be tested.

$$A(z,x) = K(x) + 4\,z^3\,B(x). \tag{3}$$

Exists ?

Laitinen, 2002



Remark 3.4. The existence theory for diffusion approximations is not entirely trivial. If conduction were not present the question could be resolved using the well-known Kirchoff transform [8,15]. In the general case more sophisticated analysis in needed. For example the methods of [10] could be tested.

$$A(z,x) = K(x) + 4\,z^3\;B(x). \tag{3}$$

Growth conditions



本章都采用求和约定，即在一项中重复脚标除特别声明外，都表示对该脚标求和。现在我们假设 a^i, b 在 $Q_T \times \boldsymbol{R} \times \boldsymbol{R}^n$ 满足结构条件：

可控增长条件：

$$|a^i(x,t,z,p)| \leqslant \mu_1(|p| + |z|^{q/n}) + \varphi_1(x,t), \qquad (1.2)$$

$$|b(x,t,z,p)| \leqslant \mu_2(|p|^{\frac{q+2}{n+2}} + |z|^{\frac{q+2}{n}}) + \varphi_2(x,t), \qquad (1.3)$$

其中 $i=1,2,\cdots,n$；

强制条件：

$$a^i(x,t,z,p)p_i \geqslant \lambda|p|^2 - |z|^2\varphi_3(x,t) - \varphi_4^2(x,t); \qquad (1.4)$$

低阶项符号条件：

$$b(x,t,z,p)z \geqslant -\lambda_0|p|^2 - |z|^2\varphi_3(x,t) - \varphi_4^2(x,t); \qquad (1.5)$$

严格单调性条件：对于任意 $(x,t,z) \in Q_T \times \boldsymbol{R}$, $p_1,p_2 \in \boldsymbol{R}^n$,

$$[a^i(x,t,z,p_1) - a^i(x,t,z,p_2)](p_{1i} - p_{2i}) > 0, \quad 当\ p_1 \neq p_2, \qquad (1.6)$$

其中 $1 \leqslant q < n+2$, $\lambda > 0$, $\lambda_0 \geqslant 0$, $\mu_i (i=1,2)$ 为非负常数，并应用了求和约定，同时还要求

$$\varphi_1 \in L^2(Q_T),\ \varphi_2 \in L^{\frac{2(n+2)}{n+4}}(Q_T),\ \varphi_3 \in L^{\frac{n+2}{2}}(Q_T),\ \varphi_4 \in L^2(Q_T).$$

Growth conditions



本章都采用求和约定，即在一项中重复脚标除特别声明外，都表示对该脚标求和．现在我们假设 a^i，b 在 $Q_T \times \mathbf{R} \times \mathbf{R}^n$ 满足**结构条件**：

可控增长条件：

$$|a^i(x,t,z,p)| \leqslant \mu_1(|p| + \boxed{|z|^{q/n}}) + \varphi_1(x,t), \qquad (1.2)$$

$$|b(x,t,z,p)| \leqslant \mu_2(|p|^{\frac{q+2}{n+2}} + \boxed{|z|^{\frac{q+2}{n}}}) + \varphi_2(x,t), \qquad (1.3)$$

其中 $i=1,2,\cdots,n$；

强制条件：

$$a^i(x,t,z,p)p_i \geqslant \lambda|p|^2 - |z|^2\varphi_3(x,t) - \varphi_4^2(x,t); \qquad (1.4)$$

低阶项符号条件：

$$b(x,t,z,p)z \geqslant -\lambda_0|p|^2 - |z|^2\varphi_3(x,t) - \varphi_4^2(x,t); \qquad (1.5)$$

严格单调性条件：对于任意 $(x,t,z) \in Q_T \times \mathbf{R}$，$p_1,p_2 \in \mathbf{R}^n$，

$$[a^i(x,t,z,p_1) - a^i(x,t,z,p_2)](p_{1i} - p_{2i}) > 0, \quad 当\ p_1 \neq p_2, \qquad (1.6)$$

其中 $\boxed{1 \leqslant q < n+2}$，$\lambda > 0, \lambda_0 \geqslant 0$，$\mu_i (i=1,2)$ 为非负常数，并应用了求和约定，同时还要求

$$\varphi_1 \in L^2(Q_T),\ \varphi_2 \in L^{\frac{2(n+2)}{n+4}}(Q_T),\ \varphi_3 \in L^{\frac{n+2}{2}}(Q_T),\ \varphi_4 \in L^2(Q_T).$$

Physical conditions

$$0 < T_{min} \leq u_b \leq T_{max}, \quad 0 \leq f \leq C.$$

$$A(z,x) = K(x) + 4\, z^3\, B(x). \tag{4}$$

If $z \in [T_{min}, T_{max}]$,

$$\lambda(T_{min}, T_{max}) \leq A(z,x) \leq \Lambda(T_{min}, T_{max}). \tag{5}$$

Physical conditions

$$0 < T_{min} \le u_b \le T_{max}, \quad 0 \le f \le C.$$

$$A(z,x) = K(x) + 4\,z^3\,B(x). \tag{4}$$

If $z \in [T_{min}, T_{max}]$,

$$\lambda(T_{min}, T_{max}) \le A(z,x) \le \Lambda(T_{min}, T_{max}). \tag{5}$$

Physical conditions

$$0 < T_{min} \le u_b \le T_{max}, \quad 0 \le f \le C.$$

$$A(z,x) = K(x) + 4\,z^3\,B(x). \tag{4}$$

If $z \in [T_{min}, T_{max}]$,

$$\lambda(T_{min}, T_{max}) \le A(z,x) \le \Lambda(T_{min}, T_{max}). \tag{5}$$

Physical conditions

$$0 < T_{min} \leq u_b \leq T_{max}, \quad 0 \leq f \leq C.$$

$$A(z,x) = K(x) + 4\, z^3\; B(x). \tag{4}$$

If $z \in [T_{min}, T_{max}]$,

$$\lambda(T_{min}, T_{max}) \leq A(z,x) \leq \Lambda(T_{min}, T_{max}). \tag{5}$$

Physical conditions

$$0 < T_{min} \leq u_b \leq T_{max}, \quad 0 \leq f \leq C.$$

$$A(z,x) = K(x) + 4\, z^3\, B(x). \tag{4}$$

If $z \in [T_{min}, T_{max}]$,

$$\lambda(T_{min}, T_{max}) \leq A(z,x) \leq \Lambda(T_{min}, T_{max}). \tag{5}$$

Physical conditions

$$0 < T_{min} \le u_b \le T_{max}, \quad 0 \le f \le C.$$

$$A(z,x) = K(x) + 4\,z^3\ B(x). \tag{4}$$

If $z \in [T_{min}, T_{max}]$,

$$\lambda(T_{min}, T_{max}) \le A(z,x) \le \Lambda(T_{min}, T_{max}). \tag{5}$$

$$\Omega \subset \mathbb{R}^n, \qquad u_b \in H^1 \cap [T_{min}, T_{max}],$$

Find $u \in H^1(\Omega)$, $\quad (u - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla \varphi = 0, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. $\quad$ Find w, $\quad (w - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \qquad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w?$

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla\varphi = 0, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla\varphi = 0, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0, \quad$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

$\Omega \subset \mathbb{R}^n, \quad u_b \in H^1 \cap [T_{min}, T_{max}],$

Find $u \in H^1(\Omega), \quad (u - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(u(x), x)\nabla u \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{6}$$

Linearized. Find $w, \quad (w - u_b)|_{\partial\Omega} = 0,$ s. t.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{7}$$

- (1) $\mathcal{L}z = w \in [T_{min},\ T_{max}]. \quad z \in [T_{min},\ T_{max}].$
- (2) $z = w$?

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.\,e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \quad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2,$ $\mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.

(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \qquad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\qquad \mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda$.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{9}$$

$\mathcal{L}z = w. \qquad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\quad \mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.\,e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda$.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{9}$$

$\mathcal{L}z = w. \quad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\quad \mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.\,e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \quad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\quad \mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ \ T_{min} \le z(x) \le T_{max},\ \ a.\,e.\ \ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda$.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \qquad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\qquad \mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \leq z(x) \leq T_{max},\ a.e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \leq A(z) \leq \Lambda$.

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \quad T_{min} \leq w \leq T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \leq C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\qquad \mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ \ T_{min} \le z(x) \le T_{max},\ \ a.e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \qquad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.\,e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \quad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ T_{min} \le z(x) \le T_{max},\ a.\,e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \quad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ \ T_{min} \leq z(x) \leq T_{max},\ \ a.\,e.\ \ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \leq A(z) \leq \Lambda.$

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \qquad T_{min} \leq w \leq T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \leq C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\qquad\mathcal{L}$ is continuous.

A fixed point.

Linearized map

$$\mathfrak{C} = \{z \in L^2(\Omega);\ \ T_{min} \le z(x) \le T_{max},\ \ a.\,e.\ in\ \Omega\}. \tag{8}$$

(1) $\mathfrak{C}$ is closed and convex in $L^2(\Omega)$.
(2) $z \in \mathfrak{C}, \quad \lambda \le A(z) \le \Lambda.$

$$\int_\Omega A({\color{red}z(x)}, x)\nabla {\color{red}w} \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{9}$$

$$\mathcal{L}z = w. \qquad T_{min} \le w \le T_{max}, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$$

(3) $\|w\|_{H^1} \le C, \quad H^1 \Subset L^2, \quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) w— A(z) —z. $\qquad \mathcal{L}$ is continuous.

A fixed point.

$A(z)$ is continuous.

$$\|z_i - z\|_2 \to 0, \quad z_i,\, z \in \mathfrak{C} \subset L^\infty\,, \qquad A(z) = K + z^3\, B,$$

$$\begin{aligned}\|A(z_i) - A(z)\|_2 &= \|(z_i - z)(z_i^2 + z_i z + z^2)\, B\|_2 \\ &\le C\|z_i - z\|_2 \to 0.\end{aligned} \tag{10}$$

$$\begin{aligned} & \int_\Omega |a_{pq}(z_i(x), x) - a_{pq}(z(x), x)|^2 \\ \le\ & \int_\Omega C|z_i(x) - z(x)|^{2\alpha} \\ \le\ & C\Big(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}}\Big)^{\alpha}\Big(\int_\Omega 1^{s'}\Big)^{\frac{1}{s'}} \\ =\ & C\Big(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}}\Big)^{\frac{1}{2}2\alpha} \\ \le\ & C\|z_i - z\|_2^{2\alpha} \to 0.\end{aligned} \tag{11}$$

$A(z)$ is continuous.

$$\|z_i - z\|_2 \to 0, \quad z_i,\, z \in \mathfrak{C} \subset L^\infty \;, \qquad A(z) = K + z^3\, B,$$

$$\begin{aligned} \|A(z_i) - A(z)\|_2 &= \|(z_i - z)(z_i^2 + z_i z + z^2)\, B\|_2 \\ &\le C\|z_i - z\|_2 \to 0. \end{aligned} \tag{10}$$

$$\begin{aligned} & \int_\Omega |a_{pq}(z_i(x), x) - a_{pq}(z(x), x)|^2 \\ \le\ & \int_\Omega C|z_i(x) - z(x)|^{2\alpha} \\ \le\ & C\Big(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}}\Big)^{\alpha}\Big(\int_\Omega 1^{s'}\Big)^{\frac{1}{s'}} \\ =\ & C\Big(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}}\Big)^{\frac{1}{2}2\alpha} \\ \le\ & C\|z_i - z\|_2^{2\alpha} \to 0. \end{aligned} \tag{11}$$

$A(z)$ is continuous.

$$\|z_i - z\|_2 \to 0, \quad z_i,\, z \in \mathfrak{C} \subset L^\infty\ , \qquad A(z) = K + z^3\, B,$$

$$\begin{aligned} \|A(z_i) - A(z)\|_2 &= \|(z_i - z)(z_i^2 + z_i z + z^2)\, B\|_2 \\ &\le C\|z_i - z\|_2 \to 0. \end{aligned} \tag{10}$$

$$\begin{aligned} & \int_\Omega |a_{pq}(z_i(x), x) - a_{pq}(z(x), x)|^2 \\ \le\ & \int_\Omega C|z_i(x) - z(x)|^{2\alpha} \\ \le\ & C(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}})^{\alpha}(\int_\Omega 1^{s'})^{\frac{1}{s'}} \\ =\ & C(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}})^{\frac{1}{2}2\alpha} \\ \le\ & C\|z_i - z\|_2^{2\alpha} \to 0. \end{aligned} \tag{11}$$

$A(z)$ is continuous.

$$\|z_i - z\|_2 \to 0, \quad z_i,\, z \in \mathfrak{C} \subset L^\infty\ , \qquad A(z) = K + z^3\, B,$$

$$\begin{aligned} \|A(z_i) - A(z)\|_2 &= \|(z_i - z)(z_i^2 + z_i z + z^2)\, B\|_2 \\ &\le C\|z_i - z\|_2 \to 0. \end{aligned} \tag{10}$$

$$\begin{aligned} & \int_\Omega |a_{pq}(z_i(x), x) - a_{pq}(z(x), x)|^2 \\ \le\ & \int_\Omega C|z_i(x) - z(x)|^{2\alpha} \\ \le\ & C(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}})^\alpha (\int_\Omega 1^{s'})^{\frac{1}{s'}} \\ =\ & C(\int_\Omega |z_i(x) - z(x)|^{\frac{2\alpha}{\alpha}})^{\frac{1}{2}2\alpha} \\ \le\ & C\|z_i - z\|_2^{2\alpha} \to 0. \end{aligned} \tag{11}$$

$\mathcal{L}$ is continuous in L^2

Suppose $z_i, z \in \mathfrak{C}$, $\|z_i - z\|_2 \to 0$, $\mathcal{L}z_i = w_i$, $\mathcal{L}z = w$.

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \le C$. $\exists\, w_0 \in H^1(\Omega)$, $(w_0 - u_b) \in H^1_0(\Omega)$,

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega; \mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0$, $\int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0$.

$w_0 = w$, $\|w_{i_k} - w_0\|_2 \to 0$, $\|w_i - w\|_2 \to 0$.

If $\|z_i - z\|_2 \to 0$, $\|w_i - w\|_2 \to 0$.

$\mathcal{L}$ is continuous in L^2

Suppose $z_i,\ z \in \mathfrak{C}$, $\quad \|z_i - z\|_2 \to 0, \quad \mathcal{L}z_i = w_i, \quad \mathcal{L}z = w.$

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \le C. \quad \exists\ w_0 \in H^1(\Omega), \quad (w_0 - u_b) \in H^1_0(\Omega),$

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega;\mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0, \qquad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0.$

$w_0 = w, \quad \|w_{i_k} - w_0\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

If $\|z_i - z\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous in L^2

Suppose $z_i,\ z \in \mathfrak{C}, \quad \|z_i - z\|_2 \to 0, \quad \mathcal{L}z_i = w_i, \quad \mathcal{L}z = w.$

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \le C. \quad \exists\ w_0 \in H^1(\Omega), \quad (w_0 - u_b) \in H^1_0(\Omega),$

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega;\mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0, \qquad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0.$

$w_0 = w, \quad \|w_{i_k} - w_0\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

If $\|z_i - z\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous in L^2

Suppose $z_i,\ z \in \mathfrak{C}, \quad \|z_i - z\|_2 \to 0, \quad \mathcal{L} z_i = w_i, \quad \mathcal{L} z = w.$

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \le C. \quad \exists\ w_0 \in H^1(\Omega), \quad (w_0 - u_b) \in H^1_0(\Omega),$

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega; \mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0, \qquad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0.$

$w_0 = w, \quad \|w_{i_k} - w_0\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

If $\|z_i - z\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous in L^2

Suppose $z_i,\ z \in \mathfrak{C}, \quad \|z_i - z\|_2 \to 0, \quad \mathcal{L} z_i = w_i, \quad \mathcal{L} z = w.$

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \leq C. \quad \exists\ w_0 \in H^1(\Omega), \quad (w_0 - u_b) \in H^1_0(\Omega),$

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega; \mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0, \qquad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0.$

$w_0 = w, \quad \|w_{i_k} - w_0\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

If $\|z_i - z\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous in L^2

Suppose $z_i,\, z \in \mathfrak{C}, \quad \|z_i - z\|_2 \to 0, \quad \mathcal{L}z_i = w_i, \quad \mathcal{L}z = w.$

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \le C. \quad \exists\ w_0 \in H^1(\Omega), \quad (w_0 - u_b) \in H^1_0(\Omega),$

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega;\, \mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0, \qquad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0.$

$w_0 = w, \quad \|w_{i_k} - w_0\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

If $\|z_i - z\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous in L^2

Suppose $z_i,\ z \in \mathfrak{C}, \quad \|z_i - z\|_2 \to 0, \quad \mathcal{L}z_i = w_i, \quad \mathcal{L}z = w.$

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{12}$$

$$\|z_i - z\|_2 \to 0, \quad \|A(z_i) - A(z)\|_2 \to 0. \tag{13}$$

$\|w_i\|_{H^1} \leq C. \quad \exists\ w_0 \in H^1(\Omega), \quad (w_0 - u_b) \in H^1_0(\Omega),$

$$\nabla w_{i_k} \rightharpoonup \nabla w_0, \quad \text{weakly in } L^2(\Omega;\mathbb{R}^n), \tag{14}$$

$$w_{i_k} \to w_0, \quad \text{strongly in } L^2(\Omega). \tag{15}$$

$\int_\Omega A(z)\nabla w_0 \cdot \nabla\varphi = 0, \qquad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0.$

$w_0 = w, \quad \|w_{i_k} - w_0\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

If $\|z_i - z\|_2 \to 0, \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous in C^0

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{16}$$

Suppose

$$\|z_i - z\|_\infty \to 0, \quad \|A(z_i) - A(z)\|_\infty \to 0, \tag{17}$$

then

$$\|w_i - w\|_{C^0(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{H^1} \to 0. \tag{18}$$

Griepentrog, Recke (2001-2010)

$$\|w_i - w\|_{C^{0,\alpha}(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{W^{1,2,\omega}} \to 0. \tag{19}$$

Mixed, parabolic systems, nonlinear.

$\mathcal{L}$ is continuous in C^0

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{16}$$

Suppose

$$\|z_i - z\|_\infty \to 0, \quad \|A(z_i) - A(z)\|_\infty \to 0, \tag{17}$$

then

$$\|w_i - w\|_{C^0(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{H^1} \to 0. \tag{18}$$

Griepentrog, Recke (2001-2010)

$$\|w_i - w\|_{C^{0,\alpha}(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{W^{1,2,\omega}} \to 0. \tag{19}$$

Mixed, parabolic systems, nonlinear.

$\mathcal{L}$ is continuous in C^0

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{16}$$

Suppose

$$\|z_i - z\|_\infty \to 0, \quad \|A(z_i) - A(z)\|_\infty \to 0, \tag{17}$$

then

$$\|w_i - w\|_{C^0(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{H^1} \to 0. \tag{18}$$

Griepentrog, Recke (2001-2010)

$$\|w_i - w\|_{C^{0,\alpha}(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{W^{1,2,\omega}} \to 0. \tag{19}$$

Mixed, parabolic systems, nonlinear.

$\mathcal{L}$ is continuous in C^0

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla \varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla \varphi = 0. \tag{16}$$

Suppose

$$\|z_i - z\|_\infty \to 0, \quad \|A(z_i) - A(z)\|_\infty \to 0, \tag{17}$$

then

$$\|w_i - w\|_{C^0(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{H^1} \to 0. \tag{18}$$

Griepentrog, Recke (2001-2010)

$$\|w_i - w\|_{C^{0,\alpha}(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{W^{1,2,\omega}} \to 0. \tag{19}$$

Mixed, parabolic systems, nonlinear.

$\mathcal{L}$ is continuous in C^0

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{16}$$

Suppose

$$\|z_i - z\|_\infty \to 0, \quad \|A(z_i) - A(z)\|_\infty \to 0, \tag{17}$$

then

$$\|w_i - w\|_{C^0(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{H^1} \to 0. \tag{18}$$

Griepentrog, Recke (2001-2010)

$$\|w_i - w\|_{C^{0,\alpha}(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{W^{1,2,\omega}} \to 0. \tag{19}$$

Mixed, parabolic systems, nonlinear.

$\mathcal{L}$ is continuous in C^0

$$\int_\Omega A(z_i)\nabla w_i \cdot \nabla\varphi = 0, \quad \int_\Omega A(z)\nabla w \cdot \nabla\varphi = 0. \tag{16}$$

Suppose

$$\|z_i - z\|_\infty \to 0, \quad \|A(z_i) - A(z)\|_\infty \to 0, \tag{17}$$

then

$$\|w_i - w\|_{C^0(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{H^1} \to 0. \tag{18}$$

Griepentrog, Recke (2001-2010)

$$\|w_i - w\|_{C^{0,\alpha}(\overline{\Omega})} \to 0, \quad \|w_i - w\|_{W^{1,2,\omega}} \to 0. \tag{19}$$

Mixed, parabolic systems, nonlinear.

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, u is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x),x)\nabla w\cdot\nabla\varphi=0,\quad \forall\ \varphi\in H^1_0(\Omega). \tag{20}$$

(1) $z\in\mathfrak{C}$, $\mathfrak{C}=\{L^2;[T_{min},T_{max}]\}$ closed convex.

(2) Maximum principles, $\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

(3) $H^1\Subset L^2$, $\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u=u$, u is a solution.

$$u\in\mathfrak{C}\cap H^1(\Omega)\cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u\cdot\overrightarrow{n}=\alpha(u-u_{gas}),\quad \alpha\in[0,C]. \tag{21}$$

$u_{gas}\in[T_{min},T_{max}]$, $\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\quad \mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad \mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, $\quad u$ is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\quad \mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad \mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, $\quad u$ is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x),x)\nabla w\cdot\nabla\varphi = 0,\quad \forall\ \varphi\in H^1_0(\Omega). \tag{20}$$

(1) $z\in\mathfrak{C}$, $\quad\mathfrak{C}=\{L^2;[T_{min},T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

(3) $H^1\Subset L^2$, $\quad\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u=u$, $\quad u$ is a solution.

$$u\in\mathfrak{C}\cap H^1(\Omega)\cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u\cdot\overrightarrow{n}=\alpha(u-u_{gas}),\quad \alpha\in[0,C]. \tag{21}$$

$u_{gas}\in[T_{min},T_{max}]$, $\quad\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x),x)\nabla w\cdot\nabla\varphi = 0, \quad \forall\,\varphi\in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\quad \mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad \mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad \mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u=u$, $\quad u$ is a solution.

$$u\in\mathfrak{C}\cap H^1(\Omega)\cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u\cdot\overrightarrow{n} = \alpha(u-u_{gas}),\quad \alpha\in[0,C]. \tag{21}$$

$u_{gas}\in[T_{min},T_{max}]$, $\quad \mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x),x)\nabla w\cdot\nabla\varphi = 0, \quad \forall\,\varphi\in H_0^1(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, u is a solution.

$$u\in\mathfrak{C}\cap H^1(\Omega)\cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u\cdot\overrightarrow{n} = \alpha(u-u_{gas}),\quad \alpha\in[0,C]. \tag{21}$$

$u_{gas}\in[T_{min},T_{max}]$, $\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x),x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}, \quad \mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, u is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}], \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\quad \mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad \mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, $\quad u$ is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x),x)\nabla w\cdot\nabla\varphi = 0, \quad \forall\, \varphi\in H_0^1(\Omega). \tag{20}$$

(1) $z\in\mathfrak{C}$, $\quad \mathfrak{C}=\{L^2;[T_{min},T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad \mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

(3) $H^1\Subset L^2$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad \mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u=u$, $\quad u$ is a solution.

$$u\in\mathfrak{C}\cap H^1(\Omega)\cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u\cdot\overrightarrow{n}=\alpha(u-u_{gas}), \quad \alpha\in[0,C]. \tag{21}$$

$u_{gas}\in[T_{min},T_{max}]$, $\quad \mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H_0^1(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\quad \mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\quad \mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, $\quad u$ is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla \varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, u is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Fixed point

$$\int_\Omega A(z(x), x)\nabla w \cdot \nabla\varphi = 0, \quad \forall\ \varphi \in H^1_0(\Omega). \tag{20}$$

(1) $z \in \mathfrak{C}$, $\mathfrak{C} = \{L^2; [T_{min}, T_{max}]\}$ closed convex.

(2) Maximum principles, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

(3) $H^1 \Subset L^2$, $\mathcal{L}\mathfrak{C}$ is precompact.

(4) The solution—data map, $\mathcal{L}$ is continuous.

A fixed point: $\mathcal{L}u = u$, u is a solution.

$$u \in \mathfrak{C} \cap H^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega}).$$

$$A\nabla u \cdot \overrightarrow{n} = \alpha(u - u_{gas}), \quad \alpha \in [0, C]. \tag{21}$$

$u_{gas} \in [T_{min}, T_{max}]$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

Heat source

$0 \leq f \leq C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$

$$0 < \lambda(T_{min}, T_{max}) \leq A(z) \leq \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$z \in \mathfrak{C} = \{L^2; \quad [\,T_{min}, \quad T_{max} + C_*\,]\,\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \leq w \leq T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \leq f \leq C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \leq f \leq C.$ $\quad \forall\, C_* > 0, \quad$ if $z \in [\, T_{min}, \quad T_{max} + C_* \,],$

$$0 < \lambda(T_{min}, T_{max}) \leq A(z) \leq \Lambda(T_{min}, T_{max}, C_*), \qquad (22)$$

$z \in \mathfrak{C} = \{L^2 ; \quad [\, T_{min}, \quad T_{max} + C_* \,]\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H^1_0(\Omega). \qquad (23)$$

$$T_{min} \leq w \leq T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \qquad (24)$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \leq f \leq C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C$. $\quad \forall\, C_* > 0$, $\quad$ if $z \in [\,T_{min}, \quad T_{max} + C_*\,]$,

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$z \in \mathfrak{C} = \{L^2 ; \quad [\,T_{min}, \quad T_{max} + C_*\,]\,\}$,

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$\lambda(T_{min}, T_{max}, C_*)$, $\quad 0 \le f \le C(T_{min}, T_{max}, C_*)$.

$C_* = C_1(C_*)$. $\quad$ Non-existence ?

Heat source

$$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$$z \in \mathfrak{C} = \{L^2; \quad [\,T_{min}, \quad T_{max} + C_*\,]\},$$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$z \in \mathfrak{C} = \{L^2; \quad [\,T_{min}, \quad T_{max} + C_*\,]\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$z \in \mathfrak{C} = \{L^2 ; \quad [\,T_{min}, \quad T_{max} + C_*\,]\,\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$z \;\in\; \mathfrak{C} \;=\; \{L^2\,; \quad [\,T_{min}, \quad T_{max} \;+\; C_*\,]\,\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{23}$$

$$T_{min} \;\le\; w \;\le\; T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\, T_{min}, \quad T_{max} + C_* \,],$$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*\,), \tag{22}$$

$$z \;\in\; \mathfrak{C} \;=\; \{L^2\,; \quad [\,T_{min}, \quad T_{max} \;+\; C_* \,]\,\},$$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \;\le\; w \;\le\; T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*), \tag{22}$$

$z \;\in\; \mathfrak{C} \;=\; \{L^2\,; \quad [\,T_{min}, \quad T_{max} \;+\; C_*\,]\,\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H^1_0(\Omega). \tag{23}$$

$$T_{min} \;\le\; w \;\le\; T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*)$. Non-existence ?

Heat source

$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*\,), \tag{22}$$

$z \in \mathfrak{C} = \{L^2 \,; \quad [\,T_{min}, \quad T_{max} + C_*\,]\,\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C. \quad \forall\, C_* > 0, \quad$ if $z \in [\, T_{min}, \quad T_{max} + C_*\,]$,

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*\,), \tag{22}$$

$z \;\in\; \mathfrak{C} \;=\; \{L^2\,; \quad [\,T_{min}, \quad T_{max} \;+\; C_*\,]\,\},$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \;\le\; w \;\le\; T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*)$.

$C_* = C_1(C_*)$. Non-existence ?

Heat source

$$0 \le f \le C. \quad \forall\, C_* > 0, \quad \text{if } z \in [\,T_{min}, \quad T_{max} + C_*\,],$$

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*\,), \tag{22}$$

$$z \in \mathfrak{C} = \{L^2 ; \quad [\,T_{min}, \quad T_{max} + C_*\,]\,\},$$

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1, \quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}.$

$\lambda(T_{min}, T_{max}, C_*), \quad 0 \le f \le C(T_{min}, T_{max}, C_*).$

$C_* = C_1(C_*).$ Non-existence ?

Heat source

$0 \le f \le C.\quad \forall\, C_* > 0,\quad$ if $z \in [\,T_{min},\quad T_{max} + C_*\,]$,

$$0 < \lambda(T_{min}, T_{max}) \le A(z) \le \Lambda(T_{min}, T_{max}, C_*\,), \tag{22}$$

$z \in \mathfrak{C} = \{L^2\,;\quad [\,T_{min},\quad T_{max} + C_*\,]\,\}$,

$$\int_\Omega A(z)\nabla w \cdot \nabla\varphi = \int_\Omega f(z)\varphi, \quad \forall\, \varphi \in H_0^1(\Omega). \tag{23}$$

$$T_{min} \le w \le T_{max} + C_1(\Omega, \lambda(T_{min}, T_{max}), \|f\|_\infty). \tag{24}$$

Let $C_* = C_1$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$\lambda(T_{min}, T_{max}, C_*)$, $\quad 0 \le f \le C(T_{min}, T_{max}, C_*)$.

$C_* = C_1(C_*)$. Non-existence ?

Parabolic model

$$S = (0, T), \quad Q_T = \Omega \times S,$$

$$\partial_t u - \operatorname{div}\left[A(u, x, t)\nabla u\right] = 0, \quad \text{in } Q_T. \tag{25}$$

$$(u - g)|_{t=0} = 0, \quad (u - g)|_{\partial\Omega} = 0. \tag{26}$$

$$g \in H^1(Q_T) \cap [T_{min}, \quad T_{max}]. \tag{27}$$

$$\lambda \leq A(z, x, t) \leq \Lambda, \quad z \in [T_{min}, \quad T_{max}]. \tag{28}$$

Parabolic model

$$S = (0, T), \quad Q_T = \Omega \times S,$$

$$\partial_t u - \operatorname{div}\left[A(u, x, t)\nabla u\right] = 0, \quad \text{in } Q_T. \tag{25}$$

$$(u - g)|_{t=0} = 0, \quad (u - g)|_{\partial\Omega} = 0. \tag{26}$$

$$g \in H^1(Q_T) \cap [T_{min}, \quad T_{max}]. \tag{27}$$

$$\lambda \leq A(z, x, t) \leq \Lambda, \quad z \in [T_{min}, \quad T_{max}]. \tag{28}$$

Parabolic model

$$S = (0, T), \quad Q_T = \Omega \times S,$$

$$\color{red}{\partial_t u} - \operatorname{div}\left[A(u,x,t)\nabla u\right] = 0, \quad \text{in } Q_T. \tag{25}$$

$$(u-g)|_{t=0} = 0, \quad (u-g)|_{\partial\Omega} = 0. \tag{26}$$

$$g \in H^1(Q_T) \cap [T_{min}, \quad T_{max}]. \tag{27}$$

$$\lambda \le A(z,x,t) \le \Lambda, \quad z \in [T_{min}, \quad T_{max}]. \tag{28}$$

Parabolic model

$$S = (0, T), \quad Q_T = \Omega \times S,$$

$$\partial_t u - \operatorname{div}\left[A(u, x, t)\nabla u\right] = 0, \quad \text{in } Q_T. \tag{25}$$

$$(u - g)|_{t=0} = 0, \quad (u - g)|_{\partial\Omega} = 0. \tag{26}$$

$$g \in H^1(Q_T) \cap [T_{min}, \quad T_{max}]. \tag{27}$$

$$\lambda \le A(z, x, t) \le \Lambda, \quad z \in [T_{min}, \quad T_{max}]. \tag{28}$$

Parabolic model

$$S = (0, T), \quad Q_T = \Omega \times S,$$

$$\partial_t u - \operatorname{div}\left[A(u, x, t)\nabla u\right] = 0, \quad \text{in } Q_T. \tag{25}$$

$$(u - g)|_{t=0} = 0, \quad (u - g)|_{\partial\Omega} = 0. \tag{26}$$

$$g \in H^1(Q_T) \cap [T_{min}, \quad T_{max}]. \tag{27}$$

$$\lambda \le A(z, x, t) \le \Lambda, \quad z \in [T_{min}, \quad T_{max}]. \tag{28}$$

Parabolic model

$$S = (0, T), \qquad Q_T = \Omega \times S,$$

$$\partial_t u - \operatorname{div}\left[A(u,x,t)\nabla u\right] = 0, \quad \text{in } Q_T. \tag{25}$$

$$(u-g)|_{t=0} = 0, \quad (u-g)|_{\partial\Omega} = 0. \tag{26}$$

$$g \in H^1(Q_T) \cap [T_{min}, \quad T_{max}]. \tag{27}$$

$$\lambda \le A(z,x,t) \le \Lambda, \quad z \in [T_{min}, \quad T_{max}]. \tag{28}$$

Weak solution

$\forall\, \varphi \in L^2(\, H_0^1),$

$$\langle \partial_t u,\, \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A(u)\nabla u \cdot \nabla \varphi = 0. \tag{29}$$

$$\text{B. C.} \quad (u-g) \in \mathcal{W} = \{v \in L^2(H_0^1);\, \partial_t v \in L^2(H^{-1})\}, \tag{30}$$

$$\mathcal{W} \hookrightarrow C([0,T];\, L^2(\Omega)), \quad \text{I. C.} \quad (u-g)(x,0) = 0. \tag{31}$$

$\langle \partial_t w,\, \varphi \rangle_{L^2(H_0^1)}$, integration by parts?

$$\begin{aligned} &\langle \partial_t w,\, \varphi \rangle_{H^1(Q_T)} \\ = \quad &[\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w\partial_t \varphi. \end{aligned} \tag{32}$$

$$w_h(x,t) \;\equiv\; \frac{1}{h}\int_t^{t+h} w(x,s)\, ds. \tag{33}$$

Weak solution

$\forall\,\varphi \in L^2(\,H_0^1),$

$$\langle \partial_t u,\, \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(u)\nabla u\cdot\nabla\varphi = 0. \tag{29}$$

$$\text{B. C.}\quad (u-g)\in\mathcal{W} = \{v\in L^2(H_0^1);\, \partial_t v\in L^2(H^{-1})\}, \tag{30}$$

$$\mathcal{W}\hookrightarrow C([0,T];\,L^2(\Omega)),\quad \text{I. C.}\quad (u-g)(x,0)=0. \tag{31}$$

$\langle \partial_t w,\, \varphi\rangle_{L^2(H_0^1)}$, integration by parts?

$$\begin{aligned}&\langle \partial_t w,\, \varphi\rangle_{H^1(Q_T)}\\ = \;& [\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w\partial_t\varphi.\end{aligned} \tag{32}$$

$$w_h(x,t) \;\equiv\; \frac{1}{h}\int_t^{t+h} w(x,s)\,ds. \tag{33}$$

Weak solution

$\forall\, \varphi \in L^2(\,H^1_0),$

$$\langle \partial_t u,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(u)\nabla u \cdot \nabla \varphi = 0. \tag{29}$$

B. C. $(u-g) \in \mathcal{W} = \{v \in L^2(H^1_0);\ \partial_t v \in L^2(H^{-1})\},$ (30)

$\mathcal{W} \hookrightarrow C([0,T];\, L^2(\Omega)),$ I. C. $(u-g)(x,0) = 0.$ (31)

$\langle \partial_t w,\, \varphi \rangle_{L^2(H^1_0)},$ integration by parts?

$$\begin{aligned} &\langle \partial_t w,\, \varphi \rangle_{H^1(Q_T)} \\ &= \ [\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w\partial_t\varphi. \end{aligned} \tag{32}$$

$$w_h(x,t) \ \equiv\ \frac{1}{h}\int_t^{t+h} w(x,s)\,ds. \tag{33}$$

Weak solution

$\forall\, \varphi \in L^2(\,H^1_0),$

$$\langle \partial_t u,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(u)\nabla u \cdot \nabla \varphi = 0. \qquad (29)$$

$$\text{B. C.} \quad (u-g) \in \mathcal{W} = \{v \in L^2(H^1_0);\, \partial_t v \in L^2(H^{-1})\}, \qquad (30)$$

$$\mathcal{W} \hookrightarrow C([0,T];\, L^2(\Omega)), \quad \text{I. C.} \quad (u-g)(x,0) = 0. \qquad (31)$$

$\langle \partial_t w,\, \varphi\rangle_{L^2(H^1_0)},$ integration by parts?

$$\langle \partial_t w,\, \varphi\rangle_{H^1(Q_T)} = [\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w\partial_t\varphi. \qquad (32)$$

$$w_h(x,t) \equiv \frac{1}{h}\int_t^{t+h} w(x,s)\,ds. \qquad (33)$$

Weak solution

$\forall\, \varphi \in L^2(\, H^1_0),$

$$\langle \partial_t u,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(u)\nabla u \cdot \nabla \varphi = 0. \tag{29}$$

$$\text{B. C.} \quad (u - g) \in \mathcal{W} = \{v \in L^2(H^1_0);\, \partial_t v \in L^2(H^{-1})\}, \tag{30}$$

$$\mathcal{W} \hookrightarrow C([0,\, T];\, L^2(\Omega)), \quad \text{I. C.} \quad (u - g)(x, 0) = 0. \tag{31}$$

$\langle \partial_t w,\, \varphi \rangle_{L^2(H^1_0)},$ integration by parts?

$$\langle \partial_t w,\, \varphi \rangle_{H^1(Q_T)}$$

$$= \quad [\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w\partial_t \varphi. \tag{32}$$

$$w_h(x, t) \equiv \frac{1}{h}\int_t^{t+h} w(x, s)\, ds. \tag{33}$$

Weak solution

$\forall\, \varphi \in L^2(\, H^1_0),$

$$\langle \partial_t u,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(u)\nabla u \cdot \nabla \varphi = 0. \tag{29}$$

$$\text{B. C.} \quad (u-g) \in \mathcal{W} = \{v \in L^2(H^1_0);\, \partial_t v \in L^2(H^{-1})\}, \tag{30}$$

$$\mathcal{W} \hookrightarrow C([0,\, T];\, L^2(\Omega)), \quad \text{I. C.} \quad (u-g)(x,0) = 0. \tag{31}$$

$\langle \partial_t w,\, \varphi \rangle_{L^2(H^1_0)},$ integration by parts?

$$\begin{aligned} &\langle \partial_t w,\, \varphi \rangle_{H^1(Q_T)} \\ = \;\; &[\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w\partial_t \varphi. \end{aligned} \tag{32}$$

$$w_h(x,t) \;\equiv\; \frac{1}{h}\int_t^{t+h} w(x,s)\,ds. \tag{33}$$

Weak solution

$\forall\, \varphi \in L^2(\, H_0^1),$

$$\langle \partial_t u,\, \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A(u)\nabla u \cdot \nabla \varphi = 0. \tag{29}$$

$$\text{B. C.} \quad (u-g) \in \mathcal{W} = \{v \in L^2(H_0^1);\, \partial_t v \in L^2(H^{-1})\}, \tag{30}$$

$$\mathcal{W} \hookrightarrow C([0,\,T];\, L^2(\Omega)), \quad \text{I. C.} \quad (u-g)(x,0) = 0. \tag{31}$$

$\langle \partial_t w,\, \varphi \rangle_{L^2(H_0^1)}$, integration by parts?

$$\begin{aligned} &\langle \partial_t w,\, \varphi \rangle_{H^1(Q_T)} \\ = \;& [\int_\Omega w\varphi]|_0^T - \iint_{Q_T} w \partial_t \varphi. \end{aligned} \tag{32}$$

$$w_h(x, t) \;\equiv\; \frac{1}{h}\int_t^{t+h} w(x,s)\,ds. \tag{33}$$

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \le \phi(x,t) \le T_{max},\ \text{a.e. in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi \rangle_{L^2(S;\,H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla \varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L}z = w.$ $\qquad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$?

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \le \phi(x,t) \le T_{max},\ \text{a.e. in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi\rangle_{L^2(S;\,H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L}z = w.$ $\qquad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$?

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \le \phi(x,t) \le T_{max},\ \text{a.e. in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi\rangle_{L^2(S;\,H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla \varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L}z = w.$ $\qquad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$?

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \leq \phi(x,t) \leq T_{max},\ \text{a.e. in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi \rangle_{L^2(S;\, H^1_0(\Omega))} + \iint_{Q_T} A(z) \nabla w \cdot \nabla \varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L} z = w.$ $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$?

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \le \phi(x,t) \le T_{max},\ \text{a. e.}\ \ \text{in}\ Q_T\}. \qquad (34)$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi \rangle_{L^2(S;\,H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi = 0. \qquad (35)$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \qquad (36)$$

$\mathcal{L}z = w. \qquad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}?$

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \leq \phi(x,t) \leq T_{max},\ \text{a.e. in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi\rangle_{L^2(S;\, H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L}z = w.$ $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}?$

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \le \phi(x,t) \le T_{max},\ \text{a.e. in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi\rangle_{L^2(S;\,H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L}z = w.$ $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}?$

Linearized map

$$\mathfrak{C} = \{\phi \in L^2(Q_T);\ T_{min} \le \phi(x,t) \le T_{max},\ \text{a. e.}\ \ \text{in } Q_T\}. \tag{34}$$

is a closed convex set in $L^2(Q_T)$.

$\forall\, \varphi \in L^2(S;\ H^1_0(\Omega))$,

$$\langle \partial_t w,\ \varphi\rangle_{L^2(S;\, H^1_0(\Omega))} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi = 0. \tag{35}$$

$$(w-g) \in \mathcal{W}, \qquad (w-g)(x,0) = 0. \tag{36}$$

$\mathcal{L}z = w. \qquad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}?$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H^1_0);\ \partial_t v \in L^2(H^{-1})\} \ = \ \overline{C^\infty([0,T];\ H^1_0(\Omega))}.$$

$$\forall\,(w-g)\in\mathcal{W},\quad \exists\,\{v_i\}\subset C^\infty(H^1_0),\quad \|w-g-v_i\|_{\mathcal{W}}\to 0,$$

$$\begin{aligned}
&\langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{L^2(H^1_0)}\\
&=\int_0^T \langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{H^1_0}\\
&=\iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+
\end{aligned}\tag{37}$$

$$\begin{aligned}
&= \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})^2_+(x,t)}{2}\\
&= \int_\Omega \frac{(v_i+g-T_{max})^2_+(x,T)}{2} - 0.
\end{aligned}\tag{38}$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H^1_0);\ \partial_t v \in L^2(H^{-1})\} = \overline{C^\infty([0,T];\ H^1_0(\Omega))}.$$

$$\forall\,(w-g) \in \mathcal{W}, \quad \exists\,\{v_i\} \subset C^\infty(H^1_0), \quad \|w-g-v_i\|_{\mathcal{W}} \to 0,$$

$$\begin{aligned}
&\langle \partial_t(v_i+g),\ (v_i+g-T_{max})_+\rangle_{L^2(H^1_0)} \\
&= \int_0^T \langle \partial_t(v_i+g),\ (v_i+g-T_{max})_+\rangle_{H^1_0} \\
&= \iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+ \qquad (37)
\end{aligned}$$

$$\begin{aligned}
&= \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})^2_+(x,t)}{2} \\
&= \int_\Omega \frac{(v_i+g-T_{max})^2_+(x,T)}{2} - 0. \qquad (38)
\end{aligned}$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H_0^1);\ \partial_t v \in L^2(H^{-1})\} \ = \ \overline{C^\infty([0,T];\ H_0^1(\Omega))}.$$

$$\forall\,(w-g) \in \mathcal{W}, \quad \exists\,\{v_i\} \subset C^\infty(H_0^1), \quad \|w-g-v_i\|_{\mathcal{W}} \to 0,$$

$$\begin{aligned}
&\langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{L^2(H_0^1)} \\
&= \int_0^T \langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{H_0^1} \\
&= \iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+ \qquad (37)
\end{aligned}$$

$$\begin{aligned}
&= \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})_+^2(x,t)}{2} \\
&= \int_\Omega \frac{(v_i+g-T_{max})_+^2(x,T)}{2} - 0. \qquad (38)
\end{aligned}$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H^1_0);\ \partial_t v \in L^2(H^{-1})\} \ = \ \overline{C^\infty([0,T];\ H^1_0(\Omega))}.$$

$$\forall\,(w-g) \in \mathcal{W}, \quad \exists\,\{v_i\} \subset C^\infty(H^1_0), \quad \|w-g-v_i\|_{\mathcal{W}} \to 0,$$

$$\begin{aligned}
&\langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{L^2(H^1_0)} \\
&= \int_0^T \langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{H^1_0} \\
&= \iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+
\end{aligned} \qquad (37)$$

$$\begin{aligned}
&= \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})^2_+(x,t)}{2} \\
&= \int_\Omega \frac{(v_i+g-T_{max})^2_+(x,T)}{2} - 0.
\end{aligned} \qquad (38)$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H_0^1);\ \partial_t v \in L^2(H^{-1})\} \;=\; \overline{C^\infty([0,T];\ H_0^1(\Omega))}.$$

$$\forall\, (w-g) \in \mathcal{W}, \quad \exists\, \{v_i\} \subset C^\infty(H_0^1), \quad \|w-g-v_i\|_{\mathcal{W}} \to 0,$$

$$\begin{aligned}
&\langle \partial_t (v_i+g),\, (v_i+g-T_{max})_+ \rangle_{L^2(H_0^1)} \\
&= \int_0^T \langle \partial_t (v_i+g),\, (v_i+g-T_{max})_+ \rangle_{H_0^1} \\
&= \iint_{Q_T} \partial_t (v_i+g) \cdot (v_i+g-T_{max})_+ \qquad (37)
\end{aligned}$$

$$\begin{aligned}
&= \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})_+^2(x,t)}{2} \\
&= \int_\Omega \frac{(v_i+g-T_{max})_+^2(x,T)}{2} - 0. \qquad (38)
\end{aligned}$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H^1_0);\ \partial_t v \in L^2(H^{-1})\} \ = \ \overline{C^\infty([0,T];\ H^1_0(\Omega))}.$$

$$\forall\,(w-g) \in \mathcal{W}, \quad \exists\,\{v_i\} \subset C^\infty(H^1_0), \quad \|w - g - v_i\|_{\mathcal{W}} \to 0,$$

$$\begin{aligned} &\langle \partial_t(v_i+g), (v_i+g-T_{max})_+\rangle_{L^2(H^1_0)} \\ &= \int_0^T \langle \partial_t(v_i+g), (v_i+g-T_{max})_+\rangle_{H^1_0} \\ &= \iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+ \end{aligned} \tag{37}$$

$$\begin{aligned} &= \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})^2_+(x,t)}{2} \\ &= \int_\Omega \frac{(v_i+g-T_{max})^2_+(x,T)}{2} - 0. \end{aligned} \tag{38}$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H_0^1);\ \partial_t v \in L^2(H^{-1})\} \;=\; \overline{C^\infty([0,T];\ H_0^1(\Omega))}.$$

$$\begin{aligned}
\forall\,(w-g)\in\mathcal{W}, \quad & \exists\,\{v_i\}\subset C^\infty(H_0^1), \quad \|w-g-v_i\|_{\mathcal{W}}\to 0, \\
& \langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{L^2(H_0^1)} \\
& = \int_0^T \langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{H_0^1} \\
& = \iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+ \qquad (37)
\end{aligned}$$

$$\begin{aligned}
& = \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})_+^2(x,t)}{2} \\
& = \int_\Omega \frac{(v_i+g-T_{max})_+^2(x,T)}{2} - 0. \qquad (38)
\end{aligned}$$

Maximum principles (1)

$$\mathcal{W} = \{v \in L^2(H_0^1);\ \partial_t v \in L^2(H^{-1})\} \ = \ \overline{C^\infty([0,T];\ H_0^1(\Omega))}.$$

$$\forall\,(w-g) \in \mathcal{W}, \quad \exists\,\{v_i\} \subset C^\infty(H_0^1), \quad \|w-g-v_i\|_{\mathcal{W}} \to 0,$$

$$\begin{aligned} &\langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{L^2(H_0^1)} \\ =& \int_0^T \langle \partial_t(v_i+g),\,(v_i+g-T_{max})_+\rangle_{H_0^1} \\ =& \iint_{Q_T} \partial_t(v_i+g)\cdot(v_i+g-T_{max})_+ \end{aligned} \qquad (37)$$

$$\begin{aligned} =& \iint_{Q_T} \partial_t \frac{(v_i+g-T_{max})_+^2(x,t)}{2} \\ =& \int_\Omega \frac{(v_i+g-T_{max})_+^2(x,T)}{2} - 0. \end{aligned} \qquad (38)$$

Maximum principles (2)

$$\langle \partial_t w, \, (w - T_{max})_+ \rangle_{L^2(H^1_0)} = \int_\Omega \frac{(w - T_{max})_+^2 (x, T)}{2} \geq 0. \tag{39}$$

$$\begin{aligned} & \iint_{Q_T} A(z) \nabla w \cdot \nabla (w - T_{max})_+ \\ = \; & \iint_{Q_T} A(z) \nabla (w - T_{max})_+ \cdot \nabla (w - T_{max})_+ \\ \geq \; & \lambda \int_S \int_\Omega |\nabla (w - T_{max})_+|^2 \\ \geq \; & C(\Omega) \lambda \int_S \int_\Omega (w - T_{max})_+^2 . \end{aligned} \tag{40}$$

Maximum principles (2)

$$\langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(H_0^1)} = \int_\Omega \frac{(w - T_{max})_+^2(x, T)}{2} \geq 0. \tag{39}$$

$$\begin{aligned} & \iint_{Q_T} A(z)\nabla w \cdot \nabla (w - T_{max})_+ \\ = & \iint_{Q_T} A(z)\nabla (w - T_{max})_+ \cdot \nabla (w - T_{max})_+ \\ \geq & \lambda \int_S \int_\Omega |\nabla (w - T_{max})_+|^2 \\ \geq & C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})_+^2. \end{aligned} \tag{40}$$

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z) \nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq \ 0 \ + \ C(\Omega) \lambda \int_S \int_\Omega (w - T_{max})_+^2 \ \geq \ 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z)\nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})_+^2 \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $\quad w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\quad \mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z) \nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})_+^2 \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z) \nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})^2_+ \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z)\nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})_+^2 \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $\qquad w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\qquad \mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z) \nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq \; 0 \; + \; C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})^2_+ \; \geq \; 0. \qquad (41) \end{aligned}$$

$w \leq T_{max}$. $\qquad w \geq T_{min}$, $\quad \mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \qquad (42)$$

$\mathcal{W} \Subset L^2(Q_T)$, $\qquad \mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H_0^1)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H_0^1(\Omega))} \\ &\qquad + \iint_{Q_T} A(z)\nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})_+^2 \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $\qquad w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H_0^1)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\qquad \mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z)\nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})^2_+ \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $\qquad w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\qquad \mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Maximum principles (3)

Let $\varphi = (w - T_{max})_+ \in L^2(H^1_0)$,

$$\begin{aligned} 0 &= \langle \partial_t w, (w - T_{max})_+ \rangle_{L^2(S;\, H^1_0(\Omega))} \\ &\qquad + \iint_{Q_T} A(z)\nabla w \cdot \nabla (w - T_{max})_+ \\ &\geq 0 + C(\Omega)\lambda \int_S \int_\Omega (w - T_{max})^2_+ \geq 0. \end{aligned} \tag{41}$$

$w \leq T_{max}$. $\qquad w \geq T_{min}$, $\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$$\|w\|^2_{\mathcal{W}} = \|w\|^2_{L^2(H^1_0)} + \|\partial_t w\|^2_{L^2(H^{-1})} \leq C. \tag{42}$$

$\mathcal{W} \Subset L^2(Q_T)$, $\qquad \mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

$\mathcal{L}$ is continuous.

$\forall\, \varphi \in L^2(H^1_0), \quad z_i,\, z \in \mathfrak{C},$

$$\langle \partial_t w_i, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z_i) \nabla w_i \cdot \nabla \varphi \;=\; 0, \qquad (43)$$

$$\langle \partial_t w, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w \cdot \nabla \varphi \;=\; 0. \qquad (44)$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\quad \partial_t w_i \rightharpoonup \partial_t w_0$, weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \quad \nabla w_i \rightharpoonup \nabla w_0$, weakly in L^2.

$$\langle \partial_t w_0, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w_0 \cdot \nabla \varphi = 0. \qquad (45)$$

$w_0 = w. \quad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous.

$$\forall\,\varphi \in L^2(H^1_0), \quad z_i,\, z \in \mathfrak{C},$$

$$\langle \partial_t w_i,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z_i) \nabla w_i \cdot \nabla \varphi \;=\; 0, \tag{43}$$

$$\langle \partial_t w,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w \cdot \nabla \varphi \;=\; 0. \tag{44}$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\quad \partial_t w_i \rightharpoonup \partial_t w_0,$ weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \quad \nabla w_i \rightharpoonup \nabla w_0,$ weakly in L^2.

$$\langle \partial_t w_0,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w_0 \cdot \nabla \varphi = 0. \tag{45}$$

$w_0 = w. \quad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous.

$\forall\, \varphi \in L^2(H^1_0), \quad z_i,\, z \in \mathfrak{C},$

$$\langle \partial_t w_i,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z_i)\nabla w_i \cdot \nabla\varphi \;=\; 0, \tag{43}$$

$$\langle \partial_t w,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi \;=\; 0. \tag{44}$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\quad \partial_t w_i \rightharpoonup \partial_t w_0,\quad$ weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \quad \nabla w_i \rightharpoonup \nabla w_0,\quad$ weakly in L^2.

$$\langle \partial_t w_0,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z)\nabla w_0 \cdot \nabla\varphi = 0. \tag{45}$$

$w_0 = w. \qquad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous.

$$\forall\, \varphi \in L^2(H^1_0), \qquad z_i,\, z \in \mathfrak{C},$$

$$\begin{aligned} \langle \partial_t w_i,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z_i) \nabla w_i \cdot \nabla \varphi &= 0, && (43) \\ \langle \partial_t w,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w \cdot \nabla \varphi &= 0. && (44) \end{aligned}$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\qquad \partial_t w_i \rightharpoonup \partial_t w_0,$ weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \qquad \nabla w_i \rightharpoonup \nabla w_0,$ weakly in L^2.

$$\langle \partial_t w_0,\, \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w_0 \cdot \nabla \varphi = 0. \qquad (45)$$

$w_0 = w. \qquad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous.

$$\forall\, \varphi \in L^2(H^1_0), \qquad z_i,\ z \in \mathfrak{C},$$

$$\begin{aligned} \langle \partial_t w_i,\ \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z_i) \nabla w_i \cdot \nabla \varphi &= 0, && (43)\\ \langle \partial_t w,\ \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w \cdot \nabla \varphi &= 0. && (44) \end{aligned}$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\quad \partial_t w_i \rightharpoonup \partial_t w_0$, weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \qquad \nabla w_i \rightharpoonup \nabla w_0$, weakly in L^2.

$$\langle \partial_t w_0,\ \varphi \rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z) \nabla w_0 \cdot \nabla \varphi = 0. \qquad (45)$$

$w_0 = w. \qquad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous.

$\forall\,\varphi \in L^2(H_0^1), \quad z_i,\, z \in \mathfrak{C},$

$$\begin{aligned} \langle \partial_t w_i,\, \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A(z_i) \nabla w_i \cdot \nabla \varphi &= 0, & (43) \\ \langle \partial_t w,\, \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A(z) \nabla w \cdot \nabla \varphi &= 0. & (44) \end{aligned}$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\quad \partial_t w_i \rightharpoonup \partial_t w_0,$ weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \quad \nabla w_i \rightharpoonup \nabla w_0,$ weakly in L^2.

$$\langle \partial_t w_0,\, \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A(z) \nabla w_0 \cdot \nabla \varphi = 0. \qquad (45)$$

$w_0 = w. \quad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

$\mathcal{L}$ is continuous.

$$\forall\, \varphi \in L^2(H^1_0), \quad z_i,\, z \in \mathfrak{C},$$

$$\langle \partial_t w_i,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z_i)\nabla w_i \cdot \nabla\varphi \;=\; 0, \tag{43}$$

$$\langle \partial_t w,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z)\nabla w \cdot \nabla\varphi \;=\; 0. \tag{44}$$

$\{w_i\}$ is bounded in $\mathcal{W}$. $\quad \partial_t w_i \rightharpoonup \partial_t w_0,$ weakly in $L^2(H^{-1})$.

$\|A(z_i) - A(z)\|_2 \to 0, \quad \nabla w_i \rightharpoonup \nabla w_0,$ weakly in L^2.

$$\langle \partial_t w_0,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(z)\nabla w_0 \cdot \nabla\varphi = 0. \tag{45}$$

$w_0 = w. \quad \mathcal{W} \Subset L^2(Q_T), \quad \|w_i - w\|_2 \to 0.$

Integral

If $u\big|_{[-h_0,0]}$ is known,

$$\partial_t w - \operatorname{div}[A(\frac{1}{h_0}\int_{t-h_0}^{t} z(x,s)\,\mathrm{d}s)\nabla w] = 0. \tag{46}$$

$$\langle \partial_t w,\, \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(\frac{1}{h_0}\int_{t-h_0}^{t} z(s)\,\mathrm{d}s)\nabla w\cdot\nabla\varphi = 0. \tag{47}$$

Similarly,

Integral

If $u|_{[-h_0,0]}$ is known,

$$\partial_t w - \mathrm{div}[A(\frac{1}{h_0}\int_{t-h_0}^{t} z(x,s)\,\mathrm{d}s)\nabla w] = 0. \tag{46}$$

$$\langle \partial_t w,\, \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(\frac{1}{h_0}\int_{t-h_0}^{t} z(s)\,\mathrm{d}s)\nabla w\cdot\nabla\varphi = 0. \tag{47}$$

Similarly,

Integral

If $u|_{[-h_0,0]}$ is known,

$$\partial_t w - \mathrm{div}[A(\frac{1}{h_0}\int_{t-h_0}^{t} z(x,s)\,\mathrm{d}s)\nabla w] = 0. \tag{46}$$

$$\langle \partial_t w,\, \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(\frac{1}{h_0}\int_{t-h_0}^{t} z(s)\,\mathrm{d}s)\nabla w\cdot\nabla\varphi = 0. \tag{47}$$

Similarly,

Integral

If $u|_{[-h_0,0]}$ is known,

$$\partial_t w - \mathrm{div}[A(\frac{1}{h_0}\int_{t-h_0}^{t} z(x,s)\,\mathrm{d}s)\nabla w] = 0. \tag{46}$$

$$\langle \partial_t w,\ \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(\frac{1}{h_0}\int_{t-h_0}^{t} z(s)\,\mathrm{d}s)\nabla w\cdot\nabla\varphi = 0. \tag{47}$$

Similarly,

Integral

If $u|_{[-h_0,0]}$ is known,

$$\partial_t w - \operatorname{div}[A(\frac{1}{h_0}\int_{t-h_0}^{t} z(x,s)\,\mathrm{d}s)\nabla w] = 0. \tag{46}$$

$$\langle \partial_t w,\ \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(\frac{1}{h_0}\int_{t-h_0}^{t} z(s)\,\mathrm{d}s)\nabla w\cdot\nabla\varphi = 0. \tag{47}$$

Similarly,

Elastic wave ...



u_{max} is independent of λ, Λ.

$$\langle w'', \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A_{ij}^{hk}(z,x,t) \frac{\partial w_i}{\partial x_k} \frac{\partial \varphi_j}{\partial x_h} = 0. \qquad (48)$$

L^∞ ?

Elastic wave ...



u_{max} is independent of λ, Λ.

$$\langle w'', \varphi \rangle_{L^2(H_0^1)} + \iint_{Q_T} A_{ij}^{hk}(z,x,t) \frac{\partial w_i}{\partial x_k} \frac{\partial \varphi_j}{\partial x_h} = 0. \qquad (48)$$

L^∞ ?

Elastic wave ...



u_{max} is independent of λ, Λ.

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A_{ij}^{hk}(z,x,t)\frac{\partial w_i}{\partial x_k}\frac{\partial \varphi_j}{\partial x_h} = 0. \qquad (48)$$

L^∞ ?

Elastic wave ...



u_{max} is independent of λ, Λ.

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A_{ij}^{hk}(z,x,t)\frac{\partial w_i}{\partial x_k}\frac{\partial \varphi_j}{\partial x_h} = 0. \qquad (48)$$

L^∞ ?

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w\cdot\nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \tag{49}$$

$$\|w\|_{C(H_0^1)}+\|w'\|_{L^\infty(L^2)}+\|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2+\|\phi_0\|_{H^1}+\|\phi_1\|_{L^2}). \tag{50}$$

Suppose (1) $f(z,x,t)\in L^2$ for any z, e.g. $f\in L^\infty$.

(2) $n=1$, $C(H_0^1)\hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t)\in[\lambda,\Lambda]$, $z\in\mathfrak{C}=\{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

$H^1(Q_T)\Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w\cdot\nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \tag{49}$$

$$\|w\|_{C(H_0^1)}+\|w'\|_{L^\infty(L^2)}+\|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2+\|\phi_0\|_{H^1}+\|\phi_1\|_{L^2}). \tag{50}$$

Suppose (1) $f(z,x,t)\in L^2$ for any z, e.g. $f\in L^\infty$.

(2) $n=1$, $C(H_0^1)\hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t)\in[\lambda,\Lambda]$, $z\in\mathfrak{C}=\{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

$H^1(Q_T)\Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w \cdot \nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \quad (49)$$

$$\|w\|_{C(H_0^1)} + \|w'\|_{L^\infty(L^2)} + \|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2 + \|\phi_0\|_{H^1} + \|\phi_1\|_{L^2}). \quad (50)$$

Suppose (1) $f(z,x,t) \in L^2$ for any z, e.g. $f \in L^\infty$.

(2) $n = 1$, $C(H_0^1) \hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t) \in [\lambda, \Lambda]$, $z \in \mathfrak{C} = \{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$H^1(Q_T) \Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w\cdot\nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \tag{49}$$

$$\|w\|_{C(H_0^1)}+\|w'\|_{L^\infty(L^2)}+\|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2+\|\phi_0\|_{H^1}+\|\phi_1\|_{L^2}). \tag{50}$$

Suppose (1) $f(z,x,t) \in L^2$ for any z, e.g. $f \in L^\infty$.

(2) $n = 1$, $C(H_0^1) \hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t) \in [\lambda, \Lambda]$, $z \in \mathfrak{C} = \{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$H^1(Q_T) \Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(x,t)\nabla w \cdot \nabla \varphi = \iint_{Q_T} f({\color{red}z}, x, t)\varphi. \quad (49)$$

$$\|w\|_{C(H^1_0)} + \|w'\|_{L^\infty(L^2)} + \|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2 + \|\phi_0\|_{H^1} + \|\phi_1\|_{L^2}). \quad (50)$$

Suppose (1) $f(z,x,t) \in L^2$ for any z, e.g. $f \in L^\infty$.

(2) $n = 1$, $C(H^1_0) \hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t) \in [\lambda, \Lambda]$, $z \in \mathfrak{C} = \{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$H^1(Q_T) \Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w\cdot\nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \quad (49)$$

$$\|w\|_{C(H_0^1)}+\|w'\|_{L^\infty(L^2)}+\|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2+\|\phi_0\|_{H^1}+\|\phi_1\|_{L^2}). \quad (50)$$

Suppose (1) $f(z,x,t)\in L^2$ for any z, e.g. $f\in L^\infty$.

(2) $n=1$, $C(H_0^1)\hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t)\in[\lambda,\Lambda]$, $z\in\mathfrak{C}=\{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

$H^1(Q_T)\Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H^1_0)} + \iint_{Q_T} A(x,t)\nabla w \cdot \nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \qquad (49)$$

$$\|w\|_{C(H^1_0)}+\|w'\|_{L^\infty(L^2)}+\|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2+\|\phi_0\|_{H^1}+\|\phi_1\|_{L^2}). \qquad (50)$$

Suppose (1) $f(z,x,t) \in L^2$ for any z, e.g. $f \in L^\infty$.

(2) $n=1$, $C(H^1_0) \hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t) \in [\lambda, \Lambda]$, $z \in \mathfrak{C} = \{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C} \subset \mathfrak{C}$.

$H^1(Q_T) \Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w\cdot\nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \quad (49)$$

$$\|w\|_{C(H_0^1)}+\|w'\|_{L^\infty(L^2)}+\|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2+\|\phi_0\|_{H^1}+\|\phi_1\|_{L^2}). \quad (50)$$

Suppose (1) $f(z,x,t)\in L^2$ for any z, e.g. $f\in L^\infty$.

(2) $n=1$, $C(H_0^1)\hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t)\in[\lambda,\Lambda]$, $z\in\mathfrak{C}=\{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

$H^1(Q_T)\Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Wave ...

$$\langle w'', \varphi\rangle_{L^2(H_0^1)} + \iint_{Q_T} A(x,t)\nabla w\cdot\nabla\varphi = \iint_{Q_T} f(z,x,t)\varphi. \tag{49}$$

$$\|w\|_{C(H_0^1)} + \|w'\|_{L^\infty(L^2)} + \|w''\|_{L^2(H^{-1})} \le C_1(\|f\|_2 + \|\phi_0\|_{H^1} + \|\phi_1\|_{L^2}). \tag{50}$$

Suppose (1) $f(z,x,t)\in L^2$ for any z, e.g. $f\in L^\infty$.

(2) $n=1$, $C(H_0^1)\hookrightarrow C^0(\overline{Q}_T)$.

(3) $A(x,t)\in[\lambda,\Lambda]$, $z\in\mathfrak{C}=\{L^2(Q_T);\ [-C_1,\ C_1]\}$.

$\mathcal{L}\mathfrak{C}\subset\mathfrak{C}$.

$H^1(Q_T)\Subset L^2(Q_T)$. $\mathcal{L}\mathfrak{C}$ is precompact.

A fixed point if $\mathcal{L}$ is continuous.

Part 2: Math algorithms

$Q_t = \Omega \times (0,t)$. If t is small enough,

$$\langle \partial_t u_{i+1}, \varphi \rangle_{W^{1,1}_D(Q_t)} + \iint_{Q_t} A(u_i) \nabla u_{i+1} \cdot \nabla \varphi + \iint_{(0,t)\times\Gamma} \alpha (u_{i+1} - u_{gas}) \varphi = \iint_{Q_t} f(u_i) \varphi. \qquad (51)$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1} - u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1}) - u(t_j)}{h}$, Rothe method.

Part 2: Math algorithms

$Q_t = \Omega \times (0,t)$. If t is small enough,

$$\langle \partial_t u_{i+1}, \varphi \rangle_{W_D^{1,1}(Q_t)} + \iint_{Q_t} A(u_i)\nabla u_{i+1}\cdot\nabla\varphi + \iint_{(0,t)\times\Gamma} \alpha(u_{i+1}-u_{gas})\varphi = \iint_{Q_t} f(u_i)\varphi. \tag{51}$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1}-u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1})-u(t_j)}{h}$, Rothe method.

Part 2: Math algorithms

$Q_t = \Omega \times (0,t)$. If t is small enough,

$$\langle \partial_t u_{i+1}, \varphi \rangle_{W_D^{1,1}(Q_t)} + \iint_{Q_t} A(u_i) \nabla u_{i+1} \cdot \nabla \varphi + \iint_{(0,t)\times\Gamma} \alpha(u_{i+1} - u_{gas})\varphi = \iint_{Q_t} f(u_i)\varphi. \tag{51}$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1} - u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1}) - u(t_j)}{h}$, Rothe method.

Part 2: Math algorithms

$Q_t = \Omega \times (0,t)$. If t is small enough,

$$\begin{aligned} \langle \partial_t u_{i+1}, \varphi \rangle_{W^{1,1}_D(Q_t)} &+ \iint_{Q_t} A(u_i) \nabla u_{i+1} \cdot \nabla \varphi \\ + \iint_{(0,t)\times\Gamma} \alpha (u_{i+1} - u_{gas}) \varphi &= \iint_{Q_t} f(u_i) \varphi. \end{aligned} \tag{51}$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1} - u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1}) - u(t_j)}{h}$, Rothe method.

Part 2: Math algorithms

$Q_t = \Omega \times (0,t)$. If t is small enough,

$$\langle \partial_t u_{i+1}, \varphi \rangle_{W^{1,1}_D(Q_t)} + \iint_{Q_t} A(u_i) \nabla u_{i+1} \cdot \nabla \varphi + \iint_{(0,t)\times\Gamma} \alpha(u_{i+1} - u_{gas})\varphi = \iint_{Q_t} f(u_i)\varphi. \tag{51}$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1} - u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1}) - u(t_j)}{h}$, Rothe method.

Part 2: Math algorithms

$Q_t = \Omega \times (0,t)$. If t is small enough,

$$\begin{aligned} &\langle \partial_t u_{i+1},\, \varphi \rangle_{W^{1,1}_D(Q_t)} + \iint_{Q_t} A(u_i)\nabla u_{i+1}\cdot\nabla\varphi \\ &+ \iint_{(0,t)\times\Gamma} \alpha(u_{i+1}-u_{gas})\varphi = \iint_{Q_t} f(u_i)\varphi. \end{aligned} \tag{51}$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1}-u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1})-u(t_j)}{h}$, Rothe method.

Part 2: Math algorithms

$Q_t = \Omega \times (0, t)$. If t is small enough,

$$\begin{aligned} \langle \partial_t u_{i+1},\, \varphi \rangle_{W^{1,1}_D(Q_t)} \;&+\; \iint_{Q_t} A(u_i)\nabla u_{i+1}\cdot\nabla\varphi \\ + \iint_{(0,t)\times\Gamma} \alpha(u_{i+1} - u_{gas})\varphi \;&=\; \iint_{Q_t} f(u_i)\varphi. \end{aligned} \tag{51}$$

$A(u_{i+1}) \approx A(u_i)$.

$A(u_{i+1}) \approx A(u_i) + A'(u_i)(u_{i+1} - u_i)$, Newton method.

$\partial_t u \approx \frac{u(t_{j+1}) - u(t_j)}{h}$, Rothe method.

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + \dots \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + \dots \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l}N_j(u_0, x, y)]dy. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j)\cdot\nabla\varphi = 0, \quad \forall\varphi \in W^1_{per}(Y). \tag{57}$$

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + \ldots \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + \ldots \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla \varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l} N_j(u_0, x, y)]dy. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j) \cdot \nabla \varphi = 0, \quad \forall \varphi \in W^1_{per}(Y). \tag{57}$$

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + \ldots \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + \ldots \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l} N_j(u_0, x, y)]dy. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j)\cdot\nabla\varphi = 0, \quad \forall\varphi \in W^1_{per}(Y). \tag{57}$$

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + \ldots \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + \ldots \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l} N_j(u_0, x, y)]dy. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j)\cdot\nabla\varphi = 0, \quad \forall\varphi \in W^1_{per}(Y). \tag{57}$$

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + ... \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + ... \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l} N_j(u_0, x, y)]dy. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j) \cdot \nabla\varphi = 0, \quad \forall\varphi \in W^1_{per}(Y). \tag{57}$$

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + \ldots \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + \ldots \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l}N_j(u_0, x, y)]\mathrm{d}y. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j)\cdot\nabla\varphi = 0, \quad \forall\varphi \in W^1_{per}(Y). \tag{57}$$

Part 3: SOTS—homogenization

$$-\mathrm{div}[A(u_\varepsilon, x, \frac{x}{\varepsilon})\nabla u_\varepsilon] = 0. \tag{52}$$

$$u_\varepsilon(x) = u_0(x) + \varepsilon u_1(x, \frac{x}{\varepsilon}) + \varepsilon^2 u_2(x, \frac{x}{\varepsilon}) + \ldots \tag{53}$$

$$A(u_\varepsilon) = A(u_0, x) + \varepsilon A_1(u_0, u_1, x, \frac{x}{\varepsilon}) + \varepsilon^2 A_2 + \ldots \tag{54}$$

$$\int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{55}$$

$$a^0_{ij}(u_0, x) = \int_Y [a_{ij}(u_0) + a_{il}(u_0)\frac{\partial}{\partial y_l}N_j(u_0, x, y)]\mathrm{d}y. \tag{56}$$

$$\int_Y A(u_0)\nabla(N_j + y_j)\cdot\nabla\varphi = 0, \quad \forall\varphi \in W^1_{per}(Y). \tag{57}$$

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\lambda \le A(z) \le \Lambda$.

The div-curl Lemma, $\lambda \le A_0(z) \le \Lambda_0$.

If $A(z)$ is continuous, $A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s.t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0$,

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0$, weakly in L^2.

Fusco, Moscariello, $A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y)$.

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\lambda \le A(z) \le \Lambda$.

The div-curl Lemma, $\lambda \le A_0(z) \le \Lambda_0$.

If $A(z)$ is continuous, $A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s.t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0$,

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0$, weakly in L^2.

Fusco, Moscariello, $A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y)$.

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\lambda \leq A(z) \leq \Lambda$.

The div-curl Lemma, $\lambda \leq A_0(z) \leq \Lambda_0$.

If $A(z)$ is continuous, $A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s. t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0$,

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0$, weakly in L^2.

Fusco, Moscariello, $A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y)$.

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\lambda \le A(z) \le \Lambda$.

The div-curl Lemma, $\lambda \le A_0(z) \le \Lambda_0$.

If $A(z)$ is continuous, $A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s.t.} \quad \int_\Omega A_0(u_0, x) \nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0$,

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0$, weakly in L^2.

Fusco, Moscariello, $A_{i\nu} \equiv A(\langle w(x) \rangle_{i\nu}, \langle x \rangle_{i\nu}, y)$.

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\lambda \leq A(z) \leq \Lambda$.

The div-curl Lemma, $\lambda \leq A_0(z) \leq \Lambda_0$.

If $A(z)$ is continuous, $A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s. t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla\varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0$,

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0$, weakly in L^2.

Fusco, Moscariello, $A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y)$.

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\qquad \lambda \le A(z) \le \Lambda.$

The div-curl Lemma, $\qquad \lambda \le A_0(z) \le \Lambda_0.$

If $A(z)$ is continuous, $\qquad A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s. t.} \quad \int_\Omega A_0(u_0, x) \nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0,$

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0, \quad$ weakly in L^2.

Fusco, Moscariello, $\quad A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y).$

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\qquad \lambda \leq A(z) \leq \Lambda.$

The div-curl Lemma, $\qquad \lambda \leq A_0(z) \leq \Lambda_0.$

If $A(z)$ is continuous, $\qquad A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s. t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0,$

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0, \quad$ weakly in L^2.

Fusco, Moscariello, $\quad A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y).$

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\qquad \lambda \leq A(z) \leq \Lambda.$

The div-curl Lemma, $\qquad \lambda \leq A_0(z) \leq \Lambda_0.$

If $A(z)$ is continuous, $\qquad A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s.t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0,$

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0,$ weakly in L^2.

Fusco, Moscariello, $\quad A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y).$

Homogenization (1)

If $z \in [T_{min}, T_{max}]$, $\lambda \le A(z) \le \Lambda$.

The div-curl Lemma, $\lambda \le A_0(z) \le \Lambda_0$.

If $A(z)$ is continuous, $A_0(z)$ is also continuous.

$$\exists u_0, \quad \text{s. t.} \quad \int_\Omega A_0(u_0, x)\nabla u_0 \cdot \nabla \varphi = 0. \tag{58}$$

$\|u_\varepsilon - u_0\|_{C^0(\overline{\Omega})} \to 0$,

$A(u_\varepsilon)\nabla u_\varepsilon \rightharpoonup A_0(u_0)\nabla u_0$, weakly in L^2.

Fusco, Moscariello, $A_{i\nu} \equiv A(\langle w(x)\rangle_{i\nu}, \langle x\rangle_{i\nu}, y)$.

Homogenization (2)

$\forall\, \varepsilon > 0,\ \exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0,\ \exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0,\ \exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0$, $\exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0,\ \exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0$, $\exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0,\ \exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall \varepsilon > 0$, $\exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall\, \varepsilon > 0,\ \exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Homogenization (2)

$\forall \varepsilon > 0$, $\exists\, u_\varepsilon$, s.t.

$$\int_\Omega A(u_\varepsilon, \frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla\varphi = 0. \qquad \|u_{\varepsilon_k} - u_0\|_{C^0} \to 0. \tag{59}$$

$$\int_\Omega A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \cdot \nabla\varphi = 0, \qquad \int_\Omega A_0(u_0)\nabla w_0 \cdot \nabla\varphi = 0. \tag{60}$$

$$\|u_{0,\varepsilon} - w_0\|_{L^2} \to 0, \quad A(u_0, \frac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla w_0. \tag{61}$$

$$\|u_\varepsilon - u_0\|_{C^0} \to 0, \quad \|A(u_\varepsilon, \tfrac{x}{\varepsilon}) - A(u_0, \tfrac{x}{\varepsilon})\|_{L^\infty} \to 0,$$

$$\|u_\varepsilon - u_{0,\varepsilon}\|_{H^1} \to 0. \quad \|u_\varepsilon - u_{0,\varepsilon}\|_{L^2} \to 0. \qquad u_0 = w_0.$$

$$A(u_\varepsilon, \tfrac{x}{\varepsilon})\nabla u_\varepsilon \approx A(u_0, \tfrac{x}{\varepsilon})\nabla u_{0,\varepsilon} \rightharpoonup A_0(u_0)\nabla u_0.$$

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y = (0,1)^n, \quad W^1_{per}(Y) = \{H^1(Y) : \text{Y-periodic}, \ \textstyle\int_Y \varphi = 0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \qquad (62)$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d = 0$.

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y = (0,1)^n, \quad W^1_{per}(Y) = \{H^1(Y) : \text{Y-periodic}, \ \textstyle\int_Y \varphi = 0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \qquad (62)$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d = 0$.

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y = (0,1)^n, \quad W^1_{per}(Y) = \{H^1(Y) : \text{Y-periodic}, \ \textstyle\int_Y \varphi = 0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall \varphi \in W^1_{per}(Y). \qquad (62)$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d = 0$.

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y = (0,1)^n, \quad W^1_{per}(Y) = \{H^1(Y) : \text{Y-periodic}, \ \textstyle\int_Y \varphi = 0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \qquad (62)$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d = 0.$

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y = (0,1)^n, \quad W^1_{per}(Y) = \{H^1(Y) : \text{Y-periodic}, \ \textstyle\int_Y \varphi = 0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \qquad (62)$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d = 0.$

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y = (0,1)^n, \quad W^1_{per}(Y) = \{H^1(Y) : \text{Y-periodic}, \ \textstyle\int_Y \varphi = 0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \tag{62}$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d = 0.$

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Part 4: Error in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$

$$Y=(0,1)^n, \quad W^1_{per}(Y)=\{H^1(Y):\text{Y-periodic}, \ \textstyle\int_Y \varphi=0\}.$$

Find $P \in W^1_{per}(Y)$, s.t.

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \qquad (62)$$

A is S P D. A, $\overrightarrow{B}$, d are Y-periodic. $\int_Y d=0.$

Lax-Milgram Lemma in $W^1_{per}(Y)$.

Translation invariance

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \qquad (63)$$

Boundary estimates ? Maximum principles ?



$$z \in \mathbb{R}^n, \quad Y_z = Y + z,$$

$$\int_{Y_z} A\nabla Q \cdot \nabla\varphi = \int_{Y_z} \overrightarrow{B} \cdot \nabla\varphi + \int_{Y_z} d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y_z). \quad (64)$$

Translation invariance

$$\int_Y A\nabla {\color{red}P}\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y). \qquad (63)$$

Boundary estimates ? Maximum principles ?



$$z\in\mathbb{R}^n,\quad Y_z=Y+z,$$

$$\int_{Y_z} A\nabla {\color{red}Q}\cdot\nabla\varphi=\int_{Y_z} \overrightarrow{B}\cdot\nabla\varphi+\int_{Y_z} d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y_z). \qquad (64)$$

Translation invariance

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y). \qquad (63)$$

Boundary estimates ? Maximum principles ?



$$z\in\mathbb{R}^n,\quad Y_z=Y+z,$$

$$\int_{Y_z} A\nabla Q\cdot\nabla\varphi=\int_{Y_z} \overrightarrow{B}\cdot\nabla\varphi+\int_{Y_z} d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y_z). \qquad (64)$$

Translation invariance

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y). \qquad (63)$$

Boundary estimates ? Maximum principles ?



$$z\in\mathbb{R}^n,\quad Y_z=Y+z,$$

$$\int_{Y_z} A\nabla Q\cdot\nabla\varphi=\int_{Y_z} \overrightarrow{B}\cdot\nabla\varphi+\int_{Y_z} d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y_z). \qquad (64)$$

Local—Global

If $\zeta \ (= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n), \quad d \in L^2_{per}(Y),$ $\int_Y d = 0,$

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \quad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If ζ $(= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n)$, $d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \quad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If $\zeta\ (= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n), \quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \quad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If ζ $(= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n)$, $\quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y)\in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K\subset\mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x)+d(x)\phi(x)=0, \quad \forall\,\phi(x)\in H^1_0(K). \tag{66}$$

If $d=0$,

$$-\mathrm{div}_x\zeta(x)=0, \quad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\quad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If $\zeta \ (= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n), \quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \quad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If $\zeta\ (= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n)$, $\quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\, \varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\, \phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \quad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$, $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If $\zeta \; (= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n), \quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\, \varphi(y) \in W^1_{per}(Y). \qquad (65)$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\, \phi(x) \in H^1_0(K). \qquad (66)$$

If $d = 0$,

$$-\mathrm{div}_x \zeta(x) = 0, \qquad \text{in } H^{-1}(K). \qquad (67)$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \qquad (68)$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$, $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If ζ $(= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n)$, $\quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y)\in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K\subset\mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x)\in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \qquad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If ζ $(= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n)$, $\quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \qquad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$; $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If $\zeta \ (= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n), \quad d \in L^2_{per}(Y),$
$\int_Y d = 0,$

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y) \in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x) \in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \qquad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi \in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$. $\quad\quad \|P\|_{W^{1,\infty}(Y)}$?

Local—Global

If ζ $(= A\nabla P - \overrightarrow{B}) \in L^2_{per}(Y;\mathbb{R}^n)$, $\quad d \in L^2_{per}(Y)$, $\int_Y d = 0$,

$$\int_Y \zeta(y)\cdot\nabla\varphi + d(y)\varphi = 0, \qquad \forall\,\varphi(y)\in W^1_{per}(Y). \tag{65}$$

For any bounded Lipschitz domain $K \subset \mathbb{R}^n$,

$$\int_K \zeta(x)\cdot\nabla\phi(x) + d(x)\phi(x) = 0, \quad \forall\,\phi(x)\in H^1_0(K). \tag{66}$$

If $d = 0$,

$$-\mathrm{div}_x\zeta(x) = 0, \qquad \text{in } H^{-1}(K). \tag{67}$$

(1986)→(1982), div-curl.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \tag{68}$$

$\|P\|_{C^{0,\alpha}(\overline{Y})}$, $\qquad \|P\|_{W^{1,\infty}(Y)}$?

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)} \le C.$ C. D. (1999).

$$u_0(x) + \varepsilon N_k\left(\frac{x}{\varepsilon}\right)\frac{\partial u_0}{\partial x_i} + \varepsilon^2 M_{kl}\left(\frac{x}{\varepsilon}\right)\frac{\partial^2 u_0}{\partial x_k\partial x_l}. \tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)}\le C.$ C. D. (1999).

$$u_0(x)+\varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i}+\varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k\partial x_l}. \tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y\overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad\forall\,\varphi\in W^1_{per}(Y).\tag{69}$$

$\|P\|_{W^{1,\infty}(Y)}\le C.$ C. D. (1999).

$$u_0(x)+\varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i}+\varepsilon^2M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2u_0}{\partial x_k\partial x_l}.\tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)} \le C.$ C. D. (1999).

$$u_0(x)+\varepsilon N_k\left(\frac{x}{\varepsilon}\right)\frac{\partial u_0}{\partial x_i}+\varepsilon^2 M_{kl}\left(\frac{x}{\varepsilon}\right)\frac{\partial^2 u_0}{\partial x_k\partial x_l}. \tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi = \int_Y \overrightarrow{B}\cdot\nabla\varphi + \int_Y d\varphi, \quad \forall\,\varphi\in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)} \le C.$ C. D. (1999).

$$u_0(x) + \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} + \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k\partial x_l}. \tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)}\le C.$ C. D. (1999).

$$u_0(x)+\varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i}+\varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k\partial x_l}. \tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P\cdot\nabla\varphi=\int_Y \overrightarrow{B}\cdot\nabla\varphi+\int_Y d\varphi,\quad \forall\,\varphi\in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)}\leq C.$ C. D. (1999).

$$u_0(x)+\varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i}+\varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k\partial x_l}. \tag{70}$$

Gradient estimates

Li and Vogelius (2000). Divergence form elliptic equations, piecewise Hölder continuous coefficients.

Global $W^{1,\infty}$ and piecewise $C^{1,\alpha}$ estimates.

$$\int_Y A\nabla P \cdot \nabla\varphi = \int_Y \overrightarrow{B} \cdot \nabla\varphi + \int_Y d\varphi, \quad \forall\, \varphi \in W^1_{per}(Y). \tag{69}$$

$\|P\|_{W^{1,\infty}(Y)} \leq C.$ C. D. (1999).

$$u_0(x) + \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} + \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{70}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \qquad (71)$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \qquad (72)$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i,\ \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. 1978}, \qquad (73)$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \qquad (74)$$

$$\left|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0\right| \le C\varepsilon; \qquad (75)$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i, \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. 1978}, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$\left|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0\right| \le C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i, \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. 1978}, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0| \le C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i, \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. 1978}, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0| \le C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i,\ \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. } 1978, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$\left|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0\right| \le C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i, \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. } 1978, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$\left|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0\right| \le C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, θ_i, $\eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \leq C\varepsilon, \quad \text{B. L. P. 1978}, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \leq C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0| \leq C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \tag{71}$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \tag{72}$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, $\theta_i,\ \eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. 1978}, \tag{73}$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \tag{74}$$

$$|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0| \le C\varepsilon; \tag{75}$$

Error estimates (1)

$$Z_\varepsilon \equiv u_\varepsilon - u_0(x) - \varepsilon N_k(\frac{x}{\varepsilon})\frac{\partial u_0}{\partial x_i} - \varepsilon^2 M_{kl}(\frac{x}{\varepsilon})\frac{\partial^2 u_0}{\partial x_k \partial x_l}. \qquad (71)$$

$$-\frac{\partial}{\partial x_i}\left(a_{ij}(\frac{x}{\varepsilon})\frac{\partial Z_\varepsilon}{\partial x_j}\right) = \varepsilon\frac{\partial \theta_i}{\partial x_i} + \varepsilon\eta, \qquad (72)$$

If $u_0 \in W^{3,q}$, piecewise Hölder continuous, θ_i, $\eta \in L^q$.

Maximum principles, De Giorgi-Nash estimates,

$$\sup_\Omega |u_\varepsilon - u_0| \le C\varepsilon, \quad \text{B. L. P. } 1978, \qquad (73)$$

$$\|u_\varepsilon - u_0 - \varepsilon u_1\|_{C^{0,\beta}(\overline{\Omega'})} \le C\varepsilon, \quad \Omega' \subset\subset \Omega; \qquad (74)$$

$$|\int_\Omega A(\frac{x}{\varepsilon})\nabla u_\varepsilon \cdot \nabla u_\varepsilon - \int_\Omega A_0 \nabla u_0 \cdot \nabla u_0| \le C\varepsilon; \qquad (75)$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\mathrm{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \le C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \le C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\mathrm{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \le C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \le C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\mathrm{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \le C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \le C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\text{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \le C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \le C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\mathrm{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \le C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \le C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\mathrm{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \le C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \le C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Error estimates (2)

Avellaneda-Lin (1987)

$$-\text{div}[A(\frac{x}{\varepsilon})\nabla u_\varepsilon] = f, \tag{76}$$

$A(y)$ is S P D, Y-periodic and $C^{0,\gamma}(\overline{Y})$ (or p-w smooth), $f \in L^q$,

$$\|\nabla u_\varepsilon\|_\infty \leq C(\|u_\varepsilon\|_\infty + \|f\|_q). \tag{77}$$

If $u_0 \in W^{4,q}_{loc}$, $A(y) \in C^{0,1}(\overline{Y})$, R.H. of $Z_\varepsilon \in L^q$,

$$\sup_{\overline{\Omega'}} |\nabla(u_\varepsilon - u_0 - \varepsilon u_1)| \leq C\varepsilon, \quad \Omega' \subset\subset \Omega. \tag{78}$$

Conclusions

- (1) PD in an interval, nonsmooth fixed point.
- (2) Mathematical algorithms.
- (3) SOTS — homogenization.
- (4) Error estimates in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$.

(a) Temperature, (b) $C^{0,\alpha}$ and p-w smooth — $W^{1,\infty}$.

Conclusions

- (1) PD in an interval, nonsmooth fixed point.
- (2) Mathematical algorithms.
- (3) SOTS — homogenization.
- (4) Error estimates in $C_{loc}^{0,\alpha}$ and $W_{loc}^{1,\infty}$.

(a) Temperature, (b) $C^{0,\alpha}$ and p-w smooth — $W^{1,\infty}$.

Conclusions

- (1) PD in an interval, nonsmooth fixed point.
- (2) Mathematical algorithms.
- (3) SOTS — homogenization.
- (4) Error estimates in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$.

(a) Temperature, (b) $C^{0,\alpha}$ and p-w smooth — $W^{1,\infty}$.

Conclusions

- (1) PD in an interval, nonsmooth fixed point.
- (2) Mathematical algorithms.
- (3) SOTS — homogenization.
- (4) Error estimates in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$.

(a) Temperature, (b) $C^{0,\alpha}$ and p-w smooth — $W^{1,\infty}$.

Conclusions

- (1) PD in an interval, nonsmooth fixed point.
- (2) Mathematical algorithms.
- (3) SOTS — homogenization.
- (4) Error estimates in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$.

(a) Temperature, (b) $C^{0,\alpha}$ and p-w smooth — $W^{1,\infty}$.

Conclusions

- (1) PD in an interval, nonsmooth fixed point.
- (2) Mathematical algorithms.
- (3) SOTS — homogenization.
- (4) Error estimates in $C^{0,\alpha}_{loc}$ and $W^{1,\infty}_{loc}$.

(a) Temperature, (b) $C^{0,\alpha}$ and p-w smooth — $W^{1,\infty}$.

Problems

- (1) Gradient estimates, Li-Vogelius (p-w), Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$, p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w), Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$, p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).

- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?

- (3) Uniqueness.

- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !

Problems

- (1) Gradient estimates, Li-Vogelius (p-w),
  Avellaneda-Lin ($A(x, \frac{x}{\varepsilon})$).
- (2) Flux estimates, $A\nabla u$, $A\nabla u_h$,
  p-w, $\|A_\varepsilon \nabla u_\varepsilon \cdot \overrightarrow{n} - A_0 \nabla u_0 \cdot \overrightarrow{n}\|_{C^0} \to 0$?
- (3) Uniqueness.
- (4) Nonlinear data–solution, L^∞ estimates for hyperbolic eq.

Thanks !